

Review: Transformations


CSE 681

Transformations

- Modeling transformations
 - build complex models by positioning (transforming) simple components relative to each other
- Viewing transformations
 - placing virtual camera in the world
 - transformation from world coordinates to camera coordinates
 - Perspective projection of 3D coordinates to 2D
- Animation
 - vary transformations over time to create motion


CSE 681

Transformations - Modeling



CSE 681


Transformations - Viewing




CSE 681

Modeling Transformations

Transform objects/points



Transform coordinate system



CSE 681

Affine Transformations

Transform $P = (x, y, z)$ to $Q = (x', y', z')$ by M .

Affine transformation:


$$\begin{aligned}x' &= m_{11}x + m_{12}y + m_{13}z + m_{14} \\y' &= m_{21}x + m_{22}y + m_{23}z + m_{24} \\z' &= m_{31}x + m_{32}y + m_{33}z + m_{34}\end{aligned}$$

CSE 681

Translation

Translation by (t_x, t_y, t_z) :

$$\begin{aligned}x' &= x + t_x \\y' &= y + t_y \\z' &= z + t_z\end{aligned}$$




CSE 681

Scaling

Scaling by (s_x, s_y, s_z) :

$$\begin{aligned}x' &= s_x x \\y' &= s_y y \\z' &= s_z z\end{aligned}$$

Uniform v. non-uniform scaling



CSE 681

Rotation

Rotation counter-clockwise by angle θ around the z-axis:

$$x' = x \cos(\theta) - y \sin(\theta)$$

$$y' = x \sin(\theta) + y \cos(\theta)$$

$$z' = z$$


Proof:

$$x = r \cos(\alpha)$$

$$y = r \sin(\alpha)$$

$$x' = r \cos(\alpha + \theta) = r \cos(\alpha) \cos(\theta) - r \sin(\alpha) \sin(\theta) = x \cos(\theta) - y \sin(\theta)$$

$$y' = r \sin(\alpha + \theta) = r \cos(\alpha) \sin(\theta) + r \sin(\alpha) \cos(\theta) = x \sin(\theta) + y \cos(\theta)$$



CSE 681


Rotation around x-axis

Rotation counter-clockwise by angle θ around the x-axis:

$$x' = x$$

$$y' = y \cos(\theta) - z \sin(\theta)$$

$$z' = y \sin(\theta) + z \cos(\theta)$$



CSE 681

Rotation around y-axis

Rotation counter-clockwise by angle θ around the y-axis:

$$y' = y$$

$$z' = z \cos(\theta) - x \sin(\theta)$$


$$x' = z \sin(\theta) + x \cos(\theta)$$

Or

$$x' = x \cos(\theta) + z \sin(\theta)$$

$$y' = y$$

$$z' = -x \sin(\theta) + z \cos(\theta)$$



CSE 681

Matrix Multiplication

Scaling by (s_x, s_y, s_z) :

$$x' = s_x x$$

$$y' = s_y y$$

$$z' = s_z z$$

$$\begin{pmatrix} x' \\ y' \\ z' \end{pmatrix} = \begin{bmatrix} s_x & 0 & 0 \\ 0 & s_y & 0 \\ 0 & 0 & s_z \end{bmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix}$$

Rotation counter-clockwise by

angle θ around the z-axis:

$$x' = x \cos(\theta) - y \sin(\theta)$$

$$y' = x \sin(\theta) + y \cos(\theta)$$

$$z' = z$$

$$\begin{pmatrix} x' \\ y' \\ z' \end{pmatrix} = \begin{bmatrix} \cos(\theta) & -\sin(\theta) & 0 \\ \sin(\theta) & \cos(\theta) & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix}$$

Translation by (t_x, t_y, t_z) :

$$x' = x + t_x$$

$$y' = y + t_y$$

$$z' = z + t_z$$

$$\begin{pmatrix} x' \\ y' \\ z' \end{pmatrix} = \begin{bmatrix} ? \\ ? \\ ? \end{bmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix}$$

CSE 681

Homogeneous Coordinates

Represent P by $(x, y, z, 1)$ and Q by $(x', y', z', 1)$.
(Homogeneous coordinates.)

Translation by (t_x, t_y, t_z) :

$$\begin{aligned}x' &= x + t_x \\y' &= y + t_y \\z' &= z + t_z\end{aligned}$$

$$\begin{pmatrix} x' \\ y' \\ z' \\ 1 \end{pmatrix} = \begin{bmatrix} 1 & 0 & 0 & t_x \\ 0 & 1 & 0 & t_y \\ 0 & 0 & 1 & t_z \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{pmatrix} x \\ y \\ z \\ 1 \end{pmatrix}$$

Scaling by (s_x, s_y, s_z) :

$$\begin{aligned}x' &= s_x x \\y' &= s_y y \\z' &= s_z z\end{aligned}$$

$$\begin{pmatrix} x' \\ y' \\ z' \\ 1 \end{pmatrix} = \begin{bmatrix} s_x & 0 & 0 & 0 \\ 0 & s_y & 0 & 0 \\ 0 & 0 & s_z & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{pmatrix} x \\ y \\ z \\ 1 \end{pmatrix}$$

CSE 681

Rotation Matrices

Rotation counter-clockwise by angle θ around the x-axis:

$$\begin{aligned}x' &= x \\y' &= y \cos(\theta) - z \sin(\theta) \\z' &= y \sin(\theta) + z \cos(\theta)\end{aligned}$$

$$\begin{pmatrix} x' \\ y' \\ z' \\ 1 \end{pmatrix} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos(\theta) & -\sin(\theta) & 0 \\ 0 & \sin(\theta) & \cos(\theta) & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{pmatrix} x \\ y \\ z \\ 1 \end{pmatrix}$$

Rotation counter-clockwise by angle θ around the y-axis:

$$\begin{aligned}x' &= x \cos(\theta) + z \sin(\theta) \\y' &= y \\z' &= -x \sin(\theta) + z \cos(\theta)\end{aligned}$$

$$\begin{pmatrix} x' \\ y' \\ z' \\ 1 \end{pmatrix} = \begin{bmatrix} \cos(\theta) & 0 & \sin(\theta) & 0 \\ 0 & 1 & 0 & 0 \\ -\sin(\theta) & 0 & \cos(\theta) & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{pmatrix} x \\ y \\ z \\ 1 \end{pmatrix}$$

Rotation counter-clockwise by angle θ around the z-axis:

$$\begin{aligned}x' &= x \cos(\theta) - y \sin(\theta) \\y' &= x \sin(\theta) + y \cos(\theta) \\z' &= z\end{aligned}$$

$$\begin{pmatrix} x' \\ y' \\ z' \\ 1 \end{pmatrix} = \begin{bmatrix} \cos(\theta) & -\sin(\theta) & 0 & 0 \\ \sin(\theta) & \cos(\theta) & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{pmatrix} x \\ y \\ z \\ 1 \end{pmatrix}$$

CSE 681

Affine Transformation Matrix

Affine transformation:

$$\begin{aligned}x' &= m_{11}x + m_{12}y + m_{13}z + m_{14} \\y' &= m_{21}x + m_{22}y + m_{23}z + m_{24} \\z' &= m_{31}x + m_{32}y + m_{33}z + m_{34}\end{aligned}$$

Transformation Matrix:

$$\begin{pmatrix} x' \\ y' \\ z' \\ 1 \end{pmatrix} = \begin{bmatrix} m_{11} & m_{12} & m_{13} & m_{14} \\ m_{21} & m_{22} & m_{23} & m_{24} \\ m_{31} & m_{32} & m_{33} & m_{34} \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{pmatrix} x \\ y \\ z \\ 1 \end{pmatrix}$$


CSE 681

Shearing

Shear along the x-axis:

$$\begin{aligned}x' &= x + hy \\y' &= y \\z' &= z\end{aligned}$$

$$\begin{pmatrix} x' \\ y' \\ z' \\ 1 \end{pmatrix} = \begin{bmatrix} 1 & h & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{pmatrix} x \\ y \\ z \\ 1 \end{pmatrix}$$



CSE 681

Reflection


Reflection across the x -axis:

$$x' = (-1)x$$

$$y' = y$$

$$z' = z$$

$$\begin{pmatrix} x' \\ y' \\ z' \\ 1 \end{pmatrix} = \begin{bmatrix} -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{pmatrix} x \\ y \\ z \\ 1 \end{pmatrix}$$



Reflection is a special case of scaling!

CSE 681

Elementary Transformations

- Translation
- Rotation
- Scaling
- Shear

What about inverses?

CSE 681

Inverse Transformations

Translation

Scale


Rotation

Shear

CSE 681

Compose Transformations


- Scale by (2, 1, 1)
- Translate by (20, 5, 0)
- Rotate by 30° counter-clockwise around the z -axis
- Translate by (0, -50, 0)



CSE 681

Compose Transformation Matrices


- Scale by (2, 1, 1);
- Translate by (20, 5, 0);
- Rotate by 30° counter-clockwise around the z-axis;
- Translate by (0, -50, 0).



$$\begin{pmatrix} x' \\ y' \\ z' \\ 1 \end{pmatrix} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & -50 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 0.86 & -0.5 & 0 & 0 \\ 0.5 & 0.86 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 & 20 \\ 0 & 1 & 0 & 5 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 2 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{pmatrix} x \\ y \\ z \\ 1 \end{pmatrix}$$

Compose Transformation Matrices

- Scale by (2, 1, 1);
- Translate by (20, 5, 0);
- Rotate by 30° counter-clockwise around the z-axis;
- Translate by (0, -50, 0).





$$\begin{pmatrix} x' \\ y' \\ z' \\ 1 \end{pmatrix} = \begin{bmatrix} 1.72 & -0.5 & 0 & 147 \\ 10 & 0.88 & 0 & 443 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{pmatrix} x \\ y \\ z \\ 1 \end{pmatrix}$$

CSE 681

Order Matters!

Transformations are not necessarily commutative!

- Translate by (20, 0, 0).
- Rotate by 30°.
- Rotate by 30°.
- Translate by (20, 0, 0).



CSE 681

Order of Transformation Matrices

Apply transformation matrices from right to left.

- Translate by (20, 0, 0).
- Rotate by 30°.

$$\begin{pmatrix} x' \\ y' \\ z' \\ 1 \end{pmatrix} = \begin{bmatrix} 0.86 & -0.5 & 0 & 0 \\ 0.5 & 0.86 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 & 20 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{pmatrix} x \\ y \\ z \\ 1 \end{pmatrix}$$

- Rotate by 30°.
- Translate by (20, 0, 0).

$$\begin{pmatrix} x' \\ y' \\ z' \\ 1 \end{pmatrix} = \begin{bmatrix} 1 & 0 & 0 & 20 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 0.86 & -0.5 & 0 & 0 \\ 0.5 & 0.86 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{pmatrix} x \\ y \\ z \\ 1 \end{pmatrix}$$

CSE 681

Which transformations commute?

- Translation and translation?
- Scaling and scaling?
- Rotation and rotation?
- Translation and scaling?
- Translation and rotation?
- Scaling and rotation?

CSE 681

Elementary Transformations

Elementary transformations:

- Translation;
- Rotation;
- Scaling;
- Shear.

Theorem: Every affine transformation can be decomposed into elementary operations.

Euler's Theorem: Every rotation around the origin can be decomposed into a rotation around the x-axis followed by a rotation around the y-axis followed by a rotation around the z-axis. (or a single rotation about an arbitrary axis).

CSE 681

Vectors

Vector $V = (V_x, V_y, V_z)$.

Homogeneous coordinates: $(V_x, V_y, V_z, 0)$.

Affine transformation:

$$\begin{pmatrix} V'_x \\ V'_y \\ V'_z \\ 0 \end{pmatrix} = \begin{bmatrix} m_{11} & m_{12} & m_{13} & m_{14} \\ m_{21} & m_{22} & m_{23} & m_{24} \\ m_{31} & m_{32} & m_{33} & m_{34} \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{pmatrix} V_x \\ V_y \\ V_z \\ 0 \end{pmatrix}$$

CSE 681

Vectors

Rotation:

$$\begin{pmatrix} V'_x \\ V'_y \\ V'_z \\ 0 \end{pmatrix} = \begin{bmatrix} \cos(\theta) & -\sin(\theta) & 0 & 0 \\ \sin(\theta) & \cos(\theta) & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{pmatrix} V_x \\ V_y \\ V_z \\ 0 \end{pmatrix}$$

Scaling:

$$\begin{pmatrix} V'_x \\ V'_y \\ V'_z \\ 0 \end{pmatrix} = \begin{bmatrix} s_x & 0 & 0 & 0 \\ 0 & s_y & 0 & 0 \\ 0 & 0 & s_z & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{pmatrix} V_x \\ V_y \\ V_z \\ 0 \end{pmatrix}$$


Translation:

(Note: No change.)

$$\begin{pmatrix} V_x \\ V_y \\ V_z \\ 0 \end{pmatrix} = \begin{bmatrix} 1 & 0 & 0 & t_x \\ 0 & 1 & 0 & t_y \\ 0 & 0 & 1 & t_z \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{pmatrix} V_x \\ V_y \\ V_z \\ 0 \end{pmatrix}$$

CSE 681

Transform Parametric Line



CSE 681

Line

transformed point on a line = point on transformed line


$$M[P(u)]^T = M[P_0 + uV]^T$$

$$M[P_0 + uV]^T = M[P_0]^T + uM[V]^T$$

Theorem: Affine transformations transform lines to lines.


CSE 681

Affine Transformation of Lines



CSE 681

Affine Combinations



CSE 681

Affine Combinations

$$M[R]^T = M[aP + bQ]^T$$

$$M[aP + bQ]^T = M[aP]^T + M[bQ]^T$$

$$M[aP]^T + M[bQ]^T = aM[P]^T + bM[Q]^T = aP' + bQ'$$

Theorem: Affine transformations preserve affine combinations.

CSE 681

Properties of Affine Transformations

- Affine transformations map lines to lines;
- Affine transformations preserve affine combinations;
- Affine transformations preserve parallelism;
- Affine transformations change volume by $|\text{Det}(M)|$;
- Any affine transformation can be decomposed into elementary transformations.
- Affine transformations [does/does not] preserve angles?
- Affine transformations [does/does not] preserve the intersection of two lines?
- Affine transformations [does/does not] preserve distances?

CSE 681

Properties of Transformation Matrices

First column is how
(1,0,0,0) transforms

$$\begin{bmatrix} ? \\ ? \\ ? \\ ? \end{bmatrix} = \begin{bmatrix} m_{11} & m_{12} & m_{13} & m_{14} \\ m_{21} & m_{22} & m_{23} & m_{24} \\ m_{31} & m_{32} & m_{33} & m_{34} \\ m_{41} & m_{42} & m_{43} & m_{44} \end{bmatrix} \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

Second column is how
(0,1,0,0) transforms

Third column is how
(0,0,1,0) transforms

Matrix holds how coordinate
axes transform and how
origin transforms

$$\begin{bmatrix} ? & ? & ? & ? \\ ? & ? & ? & ? \\ ? & ? & ? & ? \\ ? & ? & ? & ? \end{bmatrix} = \begin{bmatrix} m_{11} & m_{12} & m_{13} & m_{14} \\ m_{21} & m_{22} & m_{23} & m_{24} \\ m_{31} & m_{32} & m_{33} & m_{34} \\ m_{41} & m_{42} & m_{43} & m_{44} \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

CSE 681

Properties of Pure Rotation Matrices

Rows (columns) are orthogonal to each other


A row dot product any other row = 0

Each row (column) dot product times itself = 1

$$R^T = R^{-1}$$

CSE 681

Coordinate Frame




- Coordinate frame is given by origin ϕ and three mutually orthogonal unit vectors, i, j, k - defined in (x,y,z) space
- Mutually orthogonal (dot products): $i \cdot j = ?$; $i \cdot k = ?$; $j \cdot k = ?$.
- Unit vectors (dot products): $i \cdot i = ?$; $j \cdot j = ?$; $k \cdot k = ?$.


CSE 681

Orientation

Right handed coordinate system:




Left handed coordinate system:



CSE 681


Orientation

Right handed coordinate system:



Cross product: $i \times j = ?$

Left handed coordinate system:




Cross product: $i \times j = ?$

How do you test whether (i,j,k) is left handed or right handed?

CSE 681

Coordinate Transformations




Given object data points defined in the (i, j, k, ϕ) coordinate frame,
Given the definition of (i, j, k, ϕ) in (x, y, z) coordinates,

How do you determine the coordinates of the object data points in the $(x, y, z, 0)$ frame?

CSE 681

Coordinate change (Translation)




Change from (a,b,c,φ) coordinates to (x,y,z,0) coordinates:

1. Move (a,b,c, φ) to (x,y,z,0) and invert
2. Move data relative from (0,0,0) out to φ position by adding φ

$$\begin{pmatrix} x \\ y \\ z \\ 1 \end{pmatrix} = \begin{bmatrix} & & & \\ & & & \\ & & & \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{pmatrix} a \\ b \\ c \\ 1 \end{pmatrix}$$

CSE 681

Coordinate change (Rotation)



z-axis rotation by θ


Change from (a,b,c,φ) coordinates to (x,y,z,0) coordinates:

1. Rotate (a,b,c) by θ and invert
2. Rotate data by $-\theta$

$$\begin{pmatrix} x \\ y \\ z \\ 1 \end{pmatrix} = \begin{bmatrix} & & & \\ & & & \\ & & & \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{pmatrix} a \\ b \\ c \\ 1 \end{pmatrix}$$

CSE 681

Object Transformations



Given (i,j,k,p) defined in the (x,y,z,0) coordinate frame, Transform points defined in the (i, j, k, φ) coordinate frame to the (x,y,z,0) coordinate frame.


Rotate object to get x to line up with i, then translate to φ

$$\begin{bmatrix} 1 & 0 & 0 & \phi_x \\ 0 & 1 & 0 & \phi_y \\ 0 & 0 & 1 & \phi_z \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} i_x & j_x & k_x & 0 \\ i_y & j_y & k_y & 0 \\ i_z & j_z & k_z & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

CSE 681

Object Transformations




Affine transformation matrix:

$$\begin{bmatrix} i_x & j_x & k_x & \phi_x \\ i_y & j_y & k_y & \phi_y \\ i_z & j_z & k_z & \phi_z \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

CSE 681


Coordinate Transformations



Given (i,j,k,ϕ) defined in the $(x,y,z,0)$ coordinate frame,
 Transform points in $(i,j,k,0)$ coordinate frame to (x,y,z, ϕ) coordinate frame.
 translate by $-\phi$, then rotate to align i with x – then invert

CSE 681


Coordinate Transformations



$$R = \begin{bmatrix} i_x & i_y & i_z & 0 \\ j_x & j_y & j_z & 0 \\ k_x & k_y & k_z & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \quad T = \begin{bmatrix} 1 & 0 & 0 & -\phi_x \\ 0 & 1 & 0 & -\phi_y \\ 0 & 0 & 1 & -\phi_z \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

CSE 681

Coordinate Transformations




Apply RT to coordinate system
 Apply $(RT)^{-1}$ to data = $T^{-1}R^{-1}$

Affine transformation matrix:

$$\begin{bmatrix} i_x & j_x & k_x & \phi_x \\ i_y & j_y & k_y & \phi_y \\ i_z & j_z & k_z & \phi_z \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

CSE 681


Composition of coordinate change



M_1 changes from coordinate frame (x,y,z,θ) to (x',y',z',θ') .
 M_2 changes from coordinate frame (x',y',z',θ') to (x'',y'',z'',θ'') .
 Change from coordinate frame (x,y,z,θ) to (x'',y'',z'',θ'') ?


CSE 681

Composition of transformations - example



CSE 681


Transformations of normal vectors



- n is a unit normal vector to plane π .
 M is an affine transformation matrix.
 How is n transformed, to keep it perpendicular to the plane, under:
- translation?
 - rotation by θ ?
 - uniform scaling by s ?
 - shearing or non-uniform scaling?

CSE 681

Non-uniform scale transformation




n is a unit normal vector to plane π : $(-n_x, 0)$
 M is a non-uniform scale transformation matrix that only modifies x -coordinates
 If we just transform the n as a vector, then $M n^T = n'^T$
 But we want n'^T – that has a non-zero y -coordinate value

CSE 681

Transformations of normal vectors

Planar equation: $a x + b y + c z + d = 0$.
 Let $N = (a, b, c, d)$. (Note: (a, b, c) is normal vector)
 Let $P = (x, y, z, 1)$ be a point in plane π .
 Planar equation: $N \cdot P = 0$.
 M is an affine transformation matrix. (M^T is M transpose.)
 Let $P' = M P^T$.
 Find N' such that $N' \cdot P' = 0$ (transformed planar equation)
 $N \cdot P = 0$;
 $N P^T = 0$;
 $N (M^{-1} M) P^T = 0$;
 $(N M^{-1}) (M P^T) = 0$;
 $(N M^{-1}) P'^T = 0$;
 So $N' = N M^{-1}$
 To put in column-vector form, $(N')^T = (N M^{-1})^T = (M^{-1})^T N^T$
 So $N' = ((M^{-1})^T N^T)^T$ and $(M^{-1})^T$ is the transformation matrix to take N^T into N'^T
 Note: If M is a rotation matrix, $(M^{-1})^T = M$.



CSE 681